



Towards the probabilistic Earth-system simulator: a vision for the future of climate and weather prediction[†]

T. N. Palmer^{a,b,*}

^a*Atmospheric, Oceanic and Planetary Physics, University of Oxford, UK*

^b*European Centre for Medium-Range Weather Forecasts, Reading, UK*

*Correspondence to: T. N. Palmer, Atmospheric, Oceanic and Planetary Physics, Parks Road, Oxford OX1 3PU.
Email: t.n.palmer@atm.ox.ac.uk

[†] Based on the 2011 Royal Meteorological Society Presidential Address.

There is no more challenging problem in computational science than that of estimating, as accurately as science and technology allows, the future evolution of Earth's climate; nor indeed is there a problem whose solution has such importance and urgency. Historically, the simulation tools needed to predict climate have been developed, somewhat independently, at a number of weather and climate institutes around the world. While these simulators are individually deterministic, it is often assumed that the resulting diversity provides a useful quantification of uncertainty in global or regional predictions. However, this notion is not well founded theoretically and corresponding 'multi-simulator' estimates of uncertainty can be prone to systemic failure. Separate to this, individual institutes are now facing considerable challenges in finding the human and computational resources needed to develop more accurate weather and climate simulators with higher resolution and full Earth-system complexity. A new approach, originally designed to improve reliability in ensemble-based numerical weather prediction, is introduced to help solve these two rather different problems. Using stochastic mathematics, this approach recognizes uncertainty explicitly in the parametrized representation of unresolved climatic processes. Stochastic parametrization is shown to be more consistent with the underlying equations of motion and, moreover, provides more skilful estimates of uncertainty when compared with estimates from traditional multi-simulator ensembles, on time-scales where verification data exist. Stochastic parametrization can also help reduce long-term biases which have bedevilled numerical simulations of climate from the earliest days to the present. As a result, it is suggested that the need to maintain a large 'gene pool' of quasi-independent deterministic simulators may be obviated by the development of probabilistic Earth-system simulators. Consistent with the conclusions of the World Summit on Climate Modelling, this in turn implies that individual institutes will be able to pool human and computational resources in developing future-generation simulators, thus benefitting from economies of scale; the establishment of the Airbus consortium provides a useful analogy here. As a further stimulus for such evolution, discussion is given to a potential new synergy between the development of dynamical cores, and stochastic processing hardware. However, it is concluded that the traditional challenge in numerical weather prediction, of reducing deterministic measures of forecast error, may increasingly become an obstacle to the seamless development of probabilistic weather and climate simulators, paradoxical as that may appear at first sight. Indeed, going further, it is argued that it may be time to consider focusing operational weather forecast development entirely on high-resolution ensemble

prediction systems. Finally, by considering the exceptionally challenging problem of quantifying cloud feedback in climate change, it is argued that the development of the probabilistic Earth-system simulator may actually provide a route to reducing uncertainty in climate prediction. Copyright © 2012 Royal Meteorological Society

Key Words: Earth-system simulation; ensemble prediction; stochastic parametrization

Received 30 September 2011; Revised 30 January 2012; Accepted 14 February 2012; Published online in Wiley Online Library

Citation: Palmer TN. 2012. Towards the probabilistic Earth-system simulator: a vision for the future of climate and weather prediction. *Q. J. R. Meteorol. Soc.* DOI:10.1002/qj.1923

You can thank your lucky stars that you are not economists. Those poor souls don't even know their equations! (Sir John Mason, Director-General Meteorological Office, to his 1977 graduate intake)

I believe that the ultimate climate models ... will be stochastic, i.e., random numbers will appear somewhere in the time derivatives. (Lorenz, 1975)

1. Introduction

The problem of understanding and predicting climate is fundamentally a scientific one, but with extraordinary relevance for society. However, our understanding and ability to predict climate are still rudimentary. For example, due to profound uncertainties, primarily with the hydrological cycle, we are still unable to rule out the possibility that anthropogenic climate change will be catastrophic for humanity over the coming century, or something to which we can adapt relatively easily. Hence, while climate policy on mitigation or adaptation is rightly based on risk assessment, the risks cover a very broad range of potential outcomes, presenting a barrier to clear-cut policy and decision making. How well do we understand these uncertainties? Are they irreducible? Could the climate science community do better in reducing uncertainty? Key conclusions of this paper are that, while there are indeed irreducible uncertainties in predicting climate, and our understanding of these uncertainties is poor, new techniques promise not only to improve our ability to quantify climate prediction uncertainties more reliably, but also may actually help reduce uncertainty.

To take this further, the analysis presented in this paper suggests that development of new scientific tools to quantify uncertainty in predictions of climate more reliably, have implications for the way in which weather and climate institutes are themselves organized, both internally, and with respect to one another. For example, it could be argued that the existence of a substantial 'gene pool' of quasi-independent climate simulators^{1*} not only allows an

assessment of uncertainty in climate predictions (through the internal spread of the corresponding multi-simulator ensembles), but also engenders a spirit of competition between institutes, thereby fostering creativity. While these arguments have merits, there are counterarguments to be discussed in this paper: firstly, that multi-simulator ensembles may be prone to systemic failure due to shortcomings in the basic numerical ansatz used to formulate all contemporary simulators; and secondly that the limited human and computational resources available at the institutional level are major obstacles to the development of more accurate climate simulators.

The new scientific element introduced into this discussion hinges on a developing programme to reformulate stochastically our weather and climate prediction simulators. This 'stochastic' programme has emerged from the numerical weather prediction (NWP) community (e.g. Palmer, 1997, 2001; Buizza *et al.*, 1999), and its relevance to the climate problem can be seen as exemplifying the 'seamless prediction' philosophy (Palmer and Webster, 1993; WCRP, 2005; Slingo and Palmer, 2011) whereby the insights and constraints of NWP are brought to the climate table. The outline of the paper is as follows. In section 2, a number of reasons are given as to why incremental developments in the *status quo* for climate simulation science may not be able to provide the needed improvements in coming years. Section 3 discusses a programme to reformulate our comprehensive weather and climate simulators stochastically. Results are presented indicating how ensembles based on a single simulator with stochastic representations of simulator uncertainty can outperform the more conventional multi-simulator approach to uncertainty. Discussion of the need to integrate this stochastic approach into programmes of basic simulator development are discussed in section 4, using standard arguments familiar in other areas of physics. Section 5 discusses, briefly, a potential synergy between the development of probabilistic weather and climate simulators, and an emerging computer hardware design where exact bit reproducibility is sacrificed in order to improve energy efficiency. Section 6 presents an analysis of one obstacle to progress; indeed it is suggested that it may be time to stop production of a separate deterministic

^{1*}Throughout this paper, the word 'simulator' is used instead of the more conventional word 'model' (cf. Goldstein and Rougier, 2004). This may irritate some readers within the weather and climate community. However, for the public and many policy makers too, use of the word 'model' has a tendency to conjure up a picture of a child's toy. Some so-called climate 'sceptics' take advantage of this word association in portraying climate models merely as glorified computer games and not as the sophisticated mathematical representations of basic laws of physics that they are. When communicating with the public we have a tendency to use our own jargon, often subconsciously; hence we use the word 'model' in public because that is what we use amongst ourselves, unaware

of these pejorative word associations. Perhaps using the word 'simulator' will engender more respect for these numerical representations. Modern commercial pilots are trained almost exclusively on simulators; that apparently does not deter the public from flying. (If instead the pilots were trained merely on 'models' perhaps the public would be deterred!) As such, it may be time to start using the word 'simulator' in place of 'model' even within scientific discourse.

weather forecast, and to focus entirely on the development of probabilistic prediction systems – this may also require some evolution of practices in weather forecast offices too. Section 7 presents a vision for the development of future-generation probabilistic weather and climate simulators, using the establishment of the successful Airbus consortium as an analogy. It is argued, focusing on the thorny issue of cloud feedback in climate change prediction, that the development of the probabilistic Earth-system simulator may actually help reduce uncertainty in the magnitude (and indeed sign) of this feedback. Conclusions are given in section 8.

A key aspect of this paper is that it provides new scientific arguments to support the conclusions of the World Summit on Climate Modelling (Shukla *et al.*, 2010) that the community worldwide should be evolving towards a small number of high-resolution Earth-system simulators, possibly based the major geopolitical groupings.

Regarding the quotes at the beginning of the paper, the author was very lucky to be one of Sir John Mason's new graduate intake in 1977, and has enjoyed the most marvellous career as a result, at the Met Office, at the European Centre for Medium-Range Weather Forecasts (ECMWF), and now at Oxford University. The author agrees with Sir John's quote at the beginning of the paper, but only up to a point! And the point, as with so many other points of foundational importance on prediction and predictability, was first made by Ed Lorenz, with whom the author has had the privilege to interact during Ed's many visits to ECMWF.

In the discussion below, the importance and urgency of developing reliable climate simulators – to inform global policy on climate mitigation, to help society adapt to climate change, and to assess the impacts of proposals to actively geoengineer climate – will be assumed.

2. A critique of the traditional deterministic weather and climate simulator

2.1. The gene pool of *ab initio* climate simulators

Arrhenius (1896) developed the first mathematical simulator to quantify the effects of anthropogenic climate change. Based on the notion of energy balance in one dimension, the simulator incorporated both the direct greenhouse effect from increased carbon dioxide and the amplifying effect of water vapour, the latter through an assumption that as the atmosphere warms its relative humidity will remain constant.

The key problem with this approach is that water, unlike carbon dioxide, is not well mixed in the atmosphere, and water's three dimensional distribution, in all its phases, is sensitive to dynamical effects. The development of *ab initio* climate simulators, where dynamical effects are represented using the Navier–Stokes equations and notions such as constant relative humidity are not assumed, began with the work of Phillips (1956), who was able to adapt the simulators emerging in the rapidly developing field of NWP. The first projections of anthropogenic climate change using such *ab initio* climate simulators were given by Manabe and Wetherald (1975).

Over the years, a diversity of *ab initio* climate simulators has been produced, as individual institutes around the world sought to replicate and extend the work of these pioneers. This diversity (sometimes referred to as a 'gene pool') can

be seen as a virtue. By not putting 'all our eggs in one basket', the diversity of predictions provides an estimate of prediction uncertainty. For example, results in the IPCC Fourth Assessment Report (Solomon *et al.*, 2007) are based on a pool of coordinated projections made by some 24 climate simulators developed in different climate institutes (CMIP3; Meehl *et al.*, 2007). A similar set is currently being made for the IPCC Fifth Assessment Report.

In addition, the development of such a diversity of simulators engenders a degree of rivalry and competition between institutes that many considered necessary to foster creativity. For example, there is kudos for the institute whose climate simulator is perceived by the community as 'being the best', and having a 'world-leading' climate simulator can be considered a matter of national and institutional pride.

Maintaining such a diversity means there are relatively few opportunities to pool resources internationally and thus to benefit from 'economies of scale' when trying to improve these simulators. Hence the funding needed to improve an Earth-system simulator must largely be found at the national level. As such, even if the investment for the supercomputing needed to make global climate projections at high spatial resolution is small compared with the global costs of mitigation and adaptation, the investment may indeed be significant compared with other national funding priorities, especially in (these) times of economic difficulty.

Hence one is therefore forced to ask two questions. Notwithstanding the benefits discussed above, is this institutional-based framework unquestionably a good thing, and are the merits of the 'gene pool' incontrovertible? If not, is there an alternative?

2.2. Determinism, parametrization and scaling symmetry

All climate simulators used in CMIP3 (and indeed CMIP5) have inherited a basic feature from early NWP code: determinism². At one level, this is hardly surprising—the underlying partial differential equations on which the simulators are based (e.g. the Navier–Stokes equations) are deterministic. However, the assumption of determinism in the computational code implies that representations of unresolved processes in such simulators are themselves deterministic. For example, in his recent essay on the need for improved parametrization in atmospheric simulators, Jakob (2010) notes that, since many important processes in the atmosphere remain unresolved, 'it is therefore necessary to represent those subgrid-scale processes as a function of the grid-scale variables.' In mathematics, a function associates one quantity – the argument – with another quantity – the value – in the sense that exactly one value is associated with each argument. This characterizes perfectly the conventional approach to parametrization: the grid-scale variables determine precisely the grid-box tendency associated with the subgrid processes.

The basis for determinism appears superficially solid. Since, unlike those poor economists, we mostly know our equations at the level of partial differential equations (though see comments about Earth-system complexity near the end of section 4), we should therefore know them at the computational level too, at least at sufficiently high

²In proof, I was informed of one simulator where this is not true (Hansen *et al.*, 1983).

resolution. On top of this, improvements in deterministic parametrizations have increased the realism of comprehensive climate simulators enormously since the early days of Manabe and Wetherald, and this increase in realism has also led to substantial gains in conventional deterministic skill in weather prediction (Simmons and Hollingsworth, 2002). Is there any reason to doubt that similar improvements lie just around the corner?

However, is the argument for determinism unassailable, and is it possible that the assumption of determinism at the computational level is actually holding back progress in the development of climate and weather simulators? Let us start by going back to basics. Although the atmosphere is a compressible multi-phase fluid and indeed a considerable part of its complexity arises from this, consider for simplicity an incompressible homogeneous fluid for which the Navier–Stokes equations can be written:

$$\rho \left(\frac{\partial u}{\partial t} + u \cdot \nabla u \right) = -\nabla p + \mu \nabla^2 u, \quad (1)$$

where u is fluid velocity, p is pressure, ρ is density and μ is viscosity. These *ab initio* equations are solved numerically by truncating the equations using some finite grid or other finite (e.g. spherical harmonic) basis. If we write $u(x, t) = \bar{u}(x, t) + u'(x, t)$, where the overbar denotes some Reynolds-average operator, which we assume here to be a grid box mean, then the ‘Reynolds-averaged’ form for the Navier–Stokes equations above can be written (schematically) as

$$\rho \left(\frac{\partial \bar{u}}{\partial t} + \bar{u} \cdot \nabla \bar{u} \right) = -\nabla \bar{p} + \mu \nabla^2 \bar{u} + E$$

The effect of unresolved subgrid processes on the resolved scales are represented by the quadratic ‘Reynolds stresses’ E written in component form:

$$E_i = -\rho \nabla_j (\overline{u'_i u'_j})$$

Jakob’s definition of parametrization, applied to these Reynolds stresses, follows a long tradition in fluid dynamics, including luminaries such as Boussinesq, Prandtl, Smagorinsky (and many others), in trying to close the Reynolds-averaged equations by representing E as a deterministic function of the resolved scale variables:

$$E = P(\bar{u}; \alpha)$$

where α denotes a number of parameters which can be determined, in principle at least, by observations and/or theory.

However, a key symmetry of Eq. (1) is associated with scale invariance: if $u(x, t), p(x, t)$ is a solution to the Navier–Stokes equations, so also is

$$u_\tau(x, t) = \tau^{-1/2} u \left(\frac{x}{\tau^{1/2}}, \frac{t}{\tau} \right)$$

$$p_\tau(x, t) = \tau^{-1} p \left(\frac{x}{\tau^{1/2}}, \frac{t}{\tau} \right),$$

for any $\tau > 0$ (Majda and Bertozzi, 2001).

While we would not expect precise scale invariance of this sort to apply to the real atmosphere (not least because of latent heating and other diabatic sources), the existence




Figure 1. Variance power spectra of wind and potential temperature based on aircraft observations. The spectra of meridional wind and temperature are shifted by one and two decades to the right, respectively. Lines with slopes -3 and $-5/3$ are entered at the same relative coordinates for each variable, for comparison. From Nastrom and Gage (1985).

of such scaling symmetries in the underlying equations is consistent with observations of power-law structure in the atmosphere. Figure 1 reproduces the celebrated result of Nastrom and Gage (1985) showing an observational analysis of atmospheric kinetic energy as a function of horizontal scale (shown in terms of horizontal wave number k). This analysis draws attention to two separate power-law slopes: a ‘ -3 ’ slope at large scales and a ‘ $-5/3$ ’ slope at smaller scales. The truncation scale of all weather forecast simulators, and a number of contemporary climate simulators, lies within the ‘ $-5/3$ ’ range. Similar power-law behaviour has been seen in cloud data (Rossow and Cairns, 1995). While there is some disagreement concerning the physical interpretation of these power laws (see, for example, Lindborg, 2007), broadly speaking it appears that the ‘ -3 ’ slope is indicative of quasi-two-dimensional flow dominated by rotation, while the ‘ $-5/3$ ’ slope is indicative of three-dimensional flow with substantial divergent motion (enhanced by latent heat release in cloud systems, associated with the compressible multi-phase nature of the atmosphere).

As first clearly pointed out by Schertzer and Lovejoy (1993), the ‘deterministic truncation/parametrization ansatz’ outlined above is inconsistent with the existence of scaling symmetries and associated power-law behaviour – for the simple reason that such power laws preclude any meaningful separation between ‘resolved’ and ‘unresolved’ scales, and hence between ‘resolved’ and ‘unresolved’ processes. See also Schertzer and Lovejoy (2004). Possibly consistent with this, it can be noted that some simulators, e.g. that of ECMWF, have difficulty simulating the ‘ $-5/3$ ’ spectrum, even at relatively high truncation scales of 10 km (Straus, 2011, personal communication, using data from integrations performed as part of the Athena project; Jung *et al.*, 2012; Kinter *et al.*, 2012).

It can be argued that the failure of deterministic parametrizations to represent this observed power-law structure is the fundamental cause of systematic model error. For example, in the IPCC AR4 it is concluded that:

models still show significant errors. Although these are generally greater at smaller scales, important large-scale problems also remain. ... The ultimate source of most such errors is that many important small-scale processes cannot be represented explicitly in models, and so must be included in approximate form as they interact with larger-scale features. ... consequently models continue to display a substantial range of global temperature change in response to specified greenhouse gas forcing. (Solomon *et al.*, 2007, ch. 8)

Perhaps one could argue that with fine-enough simulator resolution large-scale errors associated with any violation of power-law behaviour can be made arbitrarily small. A simple scaling argument (Lilly, 1973; see also Palmer, 2001) indicates that this is not a reliable conclusion. Let $E(k)$ denote atmospheric kinetic energy per unit wave number, at wave number k . We can define a time-scale $\tau(k)$ in terms of a length divided by a velocity, i.e. $\tau(k) \sim k^{-\frac{3}{2}} E^{-\frac{1}{2}}(k)$. Let us suppose $\tau(k)$ characterizes the time it takes errors at wave number k to grow and infect nonlinearly the accuracy of simulations at wave number $k/2$. As above, suppose we are only interested in large-scale aspects of the flow, i.e. wave numbers less than some k_L . We can ask how long it will take before truncation errors at large wave numbers $2^N k_L, N \gg 1$ could affect large-scale simulations of the

flow. A plausible estimate of this is given by

$$\begin{aligned} \Omega(N) &= \tau(2^N k_L) + \tau(2^{N-1} k_L) + \dots + \tau(2^0 k_L) \\ &= \sum_{n=0}^N \tau(2^n k_L) \end{aligned}$$

Now if $E(k) \sim k^{-3}$ then $\tau(k)$ is independent of k and $\Omega(N)$ diverges as $N \rightarrow \infty$. This suggests that if the atmosphere was quasi two-dimensional all the way down to very small scales, errors at small scales could be ‘shielded’ from the large scales, by increasing the simulator resolution sufficiently. However, if $E(k) \sim k^{-5/3}$ then $\tau(k) \sim k^{-2/3}$ and $\Omega(N) \sim 2.7\tau(k_L)$. There is nothing especially significant about the precise value 2.7. Hence let us say that with a $-5/3$ power law the series $\Omega(N)$ converges to a value less than a few ‘eddy turnover times’ of k_L , as $N \rightarrow \infty$. Hence, with a ‘ $-5/3$ ’ power law, it may be impossible to shield the large scales from truncation-scale errors by increasing sufficiently the resolution of the simulator. This analysis is consistent with the study of Lorenz (1969); see also the more robust analysis of Rotunno and Snyder (2008) using the surface quasi-geostrophic equations, but which has not been proven rigorously from the underlying 3D Navier–Stokes equations. (It is probably not literally true in the limit where $2^N k_L \sim k_V$ and k_V lies in the viscous range of scales; however, it appears to be an open question asymptotically in the range $k_L \ll 2^N k_L \ll k_V$.) It is worth commenting that the predictability estimates above do not depend on the mechanism by which the $-5/3$ power law is established.

Despite this, there are very good reasons for attempting to increase the resolution of atmospheric simulators as much as possible. Firstly, the higher a simulator’s resolution, the better the Earth’s topography and land/sea boundary can be represented. Secondly, high resolution ensures that Rossby wave breaking, important for the maintenance of blocking anticyclones and other nonlinear weather regime phenomena (see section 7), can be simulated properly. Thirdly, the higher the resolution, the better the simulator can utilize high-resolution observations, e.g. from satellite instruments with small pixel size. Finally, at some stage, high-resolution simulators will be capable of representing the key atmospheric phenomenon of deep convection (which, along with baroclinic instability, can be considered one of the core dynamical modes of atmospheric instability and hence variability). Similar arguments apply to the oceans too. In addition to these theoretical considerations, regional predictions of climate change, particularly for precipitation change, have been shown to be sensitive to changes in resolution, horizontal and vertical (Matsueda and Palmer, 2011; Scaife *et al.*, 2011).

However, a plausible consequence of the analysis above is that as the truncation scale of a climate simulator moves into ‘ $-5/3$ ’ range the effects of the inconsistency of using deterministic parametrization cannot be reduced to zero by increasing resolution sufficiently (building a comprehensive climate simulator whose truncation scale lies in the viscous range is utterly impracticable in the foreseeable future). By this, it is not to be inferred that the effect of misrepresenting the small scales will damage the larger scales uniformly in time; that very pessimistic scenario is inconsistent with the fact that conventional NWP simulators can, from time to time at least, predict large scales very accurately, well beyond the limit $\Omega(N) \sim 2.7\tau(k_L)$. That is to say,




Figure 2. Seasonal forecast reliability diagrams for the ENSEMBLES multi-simulator ensemble. Based on 1980–2001 hindcasts initialized on 1 May and for forecast period June–August. (a) Seasonal mean NINO3 sea surface temperature above upper climatological tercile. (b) Seasonal mean precipitation anomalies in Amazon Basin in lower climatological tercile. (c) as (b) but for northern Europe. (d) As (b) but for Sahel. The dotted lines show the climatological frequency of the event and the size of the grey dots is indicative of the relative sample size within that probability bin.

experience suggests that the rapid upscale error propagation associated with the ‘ $-5/3$ ’ spectrum will occur somewhat intermittently (for example, the source of some especially erroneous medium-range weather forecasts over Europe have been traced to short-range forecast errors associated with intense mesoscale convection over the US Midwest). This raises the fundamental question: how can we ensure that the advantages of integrating simulators at higher and higher resolution will not be somehow be destroyed by rapid intermittent upscale propagation of error?

As suggested by this analysis, contemporary simulators may have common failings due to the universal use of the deterministic truncation/parametrization ansatz. This implies that multi-simulator ensembles may be blind to the consequences of such systemic failings, so that ensemble agreement cannot be assumed a reliable measure of forecast confidence. Is there any evidence for this?

There is some evidence from the poorness of the ‘attributes curve’ in reliability diagrams (Wilks, 2006) from seasonal forecasts of regional precipitation based on DEMETER multi-simulator ensembles (Palmer *et al.*, 2008). An attributes curve can assess whether, for a particular forecast event E , forecast probabilities of E are well calibrated

against observed frequencies of E – the technical definition of ‘reliability’. The attributes curve for a reliable forecast system should lie on the diagonal. Figure 2 shows an update of such seasonal forecast reliability diagrams but based on the more recent ENSEMBLES multi-simulator ensemble (Weisheimer *et al.*, 2009). Figure 2 shows examples (for seasonal mean Sahel and northern European rainfall) where the flatness of the attributes curves indicates that the ensemble is extremely overconfident and hence highly unreliable. The origin of such unreliability is, most likely, an inadequate representation of simulator error in the multi-simulator ensemble (the author is unaware of any systematic misrepresentation of observational uncertainty that would lead to such unreliability).

As discussed in Palmer *et al.* (2008), some of the unreliability of seasonal forecasts arises from difficulties which climate simulators have in simulating the statistics of weather regimes (Straus *et al.*, 2007). For example, ability to simulate anticyclonic blocking accurately is a well-known problem amongst low-resolution climate simulators. However, recent results from the Athena project (Kinter *et al.*, 2012; Jung *et al.*, 2012) suggest, even at higher resolutions, that climate simulators may have difficulty

replicating the multimodal probability distributions of regional weather regimes (Andrew Dawson, personal communication 2011), even though such multimodality is highly significant when diagnosed from reanalysis datasets. As discussed in section 7, it is suggested that an ability to simulate regional weather regimes accurately will be key to reducing uncertainty in the cloud feedback problem for predicting global climate change.

In a recent paper, Doblas-Reyes *et al.* (2011) concluded that the dominance of simulator bias in state-of-the-art coupled ocean–atmosphere simulators is a major impediment to the investigation of decadal time-scale predictability, in particular in assessing whether useful decadal predictions can be made, given our current ability to observe the subsurface ocean. Two key points can be made here. Firstly, one of the goals of the emerging programme of ‘climate services’, that of providing reliable near-term climate forecast information to a range of customers, is not likely to be met by current-generation simulators. Secondly, the value of investment in ocean (and other) observations is not being fully realized because of simulator bias. This in turn raises the following point. There have been many discussions in the community about the relative importance of funding Earth observations, *vis à vis* climate simulator development. However, this is a false dichotomy; in truth, we will only realize the full value of investment in Earth observations when climate and weather simulators are of sufficient quality to be able to ingest and utilize these observations fully (either in analysis/reanalysis mode, or in predictive mode). If the information content in an observation is being lost prematurely due to simulator bias, then the investment in producing this observation will not have been fully realized.

2.3. True diversity of the ‘gene pool’ of climate simulators

Given the problems above, it is worth asking just how diverse our ‘gene pool’ of climate simulators really is. Many climate institutes use the same basic closures in their simulators’ parametrizations; indeed some share the same parametrizations. Estimating the effective size, M_{eff} , of the CMIP3 multi-simulator ensemble has recently been studied by Pennell and Reichler (2011), who note that ‘for the full [CMIP3] 24-member ensemble, this leads to an M_{eff} that ... lies only between 7.5 and 9’. They conclude: ‘The strong similarities in model error structures found in our study indicate a considerable lack of model diversity. It is reasonable to suspect that such model similarities translate into a limited range of climate change projections.’

Hence, possibly related to the systemic problems discussed above, the effective size of the gene pool is rather small: many of the institutional simulators whose integrations are submitted to CMIP are relatively minor modifications of a small number of core simulators.

There are techniques to expand ensemble size by perturbing the parameters α within a given simulator, according to expert opinion about inherent uncertainty in the values of these parameters (Murphy *et al.*, 2004; Stainforth *et al.*, 2005; see also section 4 below). While there is certainly merit in treating these parameters as uncertain and representing this uncertainty in ‘perturbed-parameter’ ensembles, evidence to date suggests that adding perturbed-parameter integrations to a multi-simulator ensemble does not change M_{eff} by much (Masson and Knutti, 2011).

2.4. Climate complexity

Notwithstanding the remarks above, there are two fundamental problems that all climate institutes acknowledge as obstacles to the development of accurate climate simulators: insufficient human resources and insufficient computing resources. These problems are especially acute in (current) economically challenged times.

Since the days of Phillips, and Manabe and Wetherald, climate simulators have become increasingly complex. In terms of parametrizations, the subgrid representations for deep convection, clear-sky and cloud radiative effects, subgrid orography, boundary layer turbulence, aerosols, cloud microphysics, etc., have become immeasurably more sophisticated (and computationally demanding) since the early days. Moreover, what in the 1970s were essentially atmosphere-only simulators (e.g. with simple ‘slab’ oceans and ‘bucket’ land–surface hydrology) have in the 2010s become fully coupled representations of the atmosphere, oceans, cryosphere and land surface with a range of biogeochemical processes (‘Earth-system complexity’). The need to ensure that chemical tracers are properly represented during simulations, yet at the same time allowing the simulators to run efficiently on massively parallel computers, means that the numerics of the dynamical cores of weather and climate simulators have to be extremely sophisticated.

Problems of algorithmic complexity do not stop there. For climate-service applications, shorter-range decadal predictions require that simulators are initialized with contemporary observations, implying the need for sophisticated data assimilation schemes for the atmosphere, oceans and land surface.

Finally, the dynamical cores themselves are increasingly complex as quasi-geostrophic equations have given rise to the hydrostatic primitive equations, and now to the non-hydrostatic dynamical cores, needed to be able to probe kilometre truncation scales where deep convection is at least partially resolved. At these high resolutions, it is a highly non-trivial problem to ensure that numerical code can run efficiently over the very large numbers of processors of modern supercomputers (the scalability problem).

Not surprisingly, climate institutes struggle to find the human resources needed to develop these manifold elements. On top of this, the computational demands of a contemporary climate simulator means it is impossible for an institute to develop simulators both with full Earth-system complexity and with the resolution of a contemporary NWP simulator, and at the same time run large ensemble integrations from states initialized with contemporary observations. This extremely important issue has been discussed at length elsewhere, being a key topic of the major World Summit on Climate Modelling (Shukla *et al.*, 2010; Palmer, 2011).

In the next sections, we discuss a relatively new approach to the representation of unresolved processes in weather and climate simulators, which may provide a solution to the complex and challenging problems outlined in this section.

3. Stochastic representation of unresolved processes

Let us begin by considering a generalization of the definition of what we mean by ‘parametrization’ and frame it, not in terms of functional relationships, but as a constraint on some prior (e.g. climatological) probability distribution of subgrid

tendency based on a knowledge of contemporaneous values of grid-scale variables. An explicit example will be given below. This automatically suggests we treat the notion of parametrization as an inherently probabilistic problem, to be tackled by explicitly stochastic techniques (Palmer, 2001).

There is nothing new in the use of stochastic mathematics to describe climate simulators; the idea can be traced to Hasselmann (1976), who developed an idealized coupled ocean–atmosphere simulator in which the entire atmosphere was represented by a simple Markov process. Using this simulator, Hasselmann showed how ocean–atmosphere coupling would redden the spectrum of atmospheric variability. However, the use of stochastic mathematics in such earlier approaches is conceptually different from the concept being explored here: Hasselmann’s

simulator is (deliberately) a simplified idealized representation of climate, and the use of stochastic mathematics made the representation of internal atmospheric variability in the simulator equations mathematically tractable. Here, we are not interested in mathematical tractability *per se*. Rather it is being argued that stochastic mathematics also has an inherent role to play in comprehensive *ab initio* weather and climate simulators.

A key conceptual difference between deterministic and stochastic parametrization is illustrated in Figure 3. While deterministic parametrization represents the bulk-average effect of some putative large ensemble of subgrid processes occurring on scales smaller than the grid scale, stochastic parametrization attempts to represent actual realizations of the subgrid flow when no scale separation exists.




Figure 3. (a) Schematic of hypothetical situation where there is some scale separation between resolved and unresolved flow, justifying the notion of deterministic parametrization. (b) Schematic of the more realistic situation where there is no scale separation between resolved and unresolved flow, justifying the notion of stochastic parametrization.

Figure 3 indicates that the stochastic parametrizations must necessarily impact directly on scales larger than the truncation scale. This is because, as discussed above, with power-law behaviour, uncertainty in subgrid processes will propagate upscale by nonlinear dynamical effects (Thuburn *et al.*, 2011). Hence part of the (stochastic) parametrization process requires one to represent the effect of uncertainty in the subgrid processes on the resolved grid.

In order to quantify the potential benefits of this stochastic approach to parametrization, it is useful to consider a reasonably tractable example where we know precisely the ‘true’ system, which we will attempt to simulate approximately using parametrizations, both deterministic and stochastic. Consider, then, the set of linked nonlinear ordinary differential equations put forward by Lorenz (1996):

$$\frac{dX_k}{dt} = -X_{k-1}(X_{k-2} - X_{k+1}) - X_k + F - \frac{hc}{b} \sum_{j=(k-1)+1}^{kj} Y_j, \quad (2)$$

$$\frac{dY_j}{dt} = -cbY_{j+1}(Y_{j+2} - Y_{j-1}) - cY_j + \frac{hc}{b} X_{int[(j-1)/J]+1}. \quad (3)$$

Here the X_k represent the large, slow scales (analogous to wave numbers $\leq k_L$) that we are interested in, and the Y_j represent the small, fast scales (analogous to wave numbers $\geq 2^N k_L$) that we wish to parametrize. Here $1 \leq j \leq 32$, and k is cyclic mod 8. The last term of the first equation couples the small scales to the large scales; we will call this ‘the small-scale tendency’. Below we consider two values of the c parameter: $c = 10$ and $c = 4$; the h, b and F parameters are held fixed. When $c = 10$, the Y variables typically evolve over substantially faster time-scales than do the X variables, i.e. there is clear temporal scale separation between these variables. It will turn out that parametrizing the small scales deterministically will work reasonably well for this parameter setting. By contrast, when $c = 4$, this scale separation is weaker and the parametrization problem becomes inherently less deterministic. By way of analogy, then, we use the values $c = 10$ to mimic the relatively steep ‘ -3 ’ energy spectrum, and $c = 4$ to represent the relatively shallow ‘ $-5/3$ ’ energy spectrum of the real atmosphere.

With the true system represented by Eqs (2) and (3), we now consider a simulator

$$\begin{aligned} \frac{dX_k}{dt} &= -X_{k-1}(X_{k-2} - X_{k+1}) - X_k + F - P_k, \\ P_k &= (1 + r_k^{\text{mult}})P_k^{\text{det}}(X_k; \alpha) + r_k^{\text{add}}, \end{aligned}$$

of the ‘true’ Lorenz (1996) system, where the small-scale tendency is parametrized by the formulae P_k (first discussed by Wilks, 2005). Here we have generalized the conventional deterministic formula $P_k = P_k^{\text{det}}(X_k; \alpha)$ using stochastic variables r_k^{add} and r_k^{mult} . A number of parametrizations are considered: ‘deterministic’ denotes a deterministic parametrization ($r_k^{\text{add}} = r_k^{\text{mult}} = 0$) based on fitting a cubic polynomial in X_k to points in a scatter diagram of instantaneous small-scale tendency against X_k ; ‘white additive’ denotes a simple white-noise term added to the deterministic parametrization ($r_k^{\text{add}} \neq 0$; $r_k^{\text{mult}} = 0$); ‘red additive’ denotes a red-noise AR1 process added to

the deterministic parametrization; ‘multiplicative’ denotes a red-noise AR1 process multiplying the tendencies from the deterministic parametrization ($r_k^{\text{add}} = 0$; $r_k^{\text{mult}} \neq 0$).

We can use this system to illustrate the utility of the probabilistic notion of parametrization as defined earlier in this section. Figure 4 shows (solid curve) the unconstrained (i.e. climatological) probability distribution of the small-scale tendency term, the last term on the right-hand side of Eq. (2) when $c = 4$. In this figure is plotted the probability distribution of this tendency when the X_k variable is constrained to lie in $-6 \leq X_k \leq -5$ (dotted line) and $13 \leq X_k \leq 14$ (dashed line). It can be seen that the constrained probability distributions are quite different from the climatological distribution. That is, knowledge of the large-scale variable is important in constraining the prior distribution. However, this knowledge does not constrain the distribution so much that it collapses to a Dirac delta function – which would be the case if deterministic parametrization were accurate. Corresponding hat functions for the putative deterministic parametrization, for $-6 \leq X_k \leq -5$ and $13 \leq X_k \leq 14$, are shown in Figure 4 for $c = 4$; compared with the constrained probability distributions, these hat functions are quite obviously too sharp. As such, it can be expected that the simulator with deterministic parametrization will perform relatively poorly. Figure 4 also shows that the probability distributions are sharper for small deterministic tendency, suggesting that the simulator with multiplicative noise parametrization may be especially skilful.

Figure 5 shows skill score results for a large number of initial-value ensemble predictions (Figure 5(a)) and one long climate integration (Figure 5(b)). Full details are given in




Figure 4. Probability distributions of the tendency term in the (‘large-scale’) X equations, due to the (‘small-scale’) Y variables in the Lorenz (1996) dynamical system with $c = 4$. Thick solid line, prior climatological distribution; dashed line, distribution conditioned on $-6 < X < -5$; dotted line, distribution conditioned on $13 < X < 14$. The fact that the distribution is broader when X is constrained to large values than when constrained to small values provides some explanation for why the multiplicative noise parametrization is more skilful in Figure 5. The thin solid lines define hat functions associated with the deterministic parametrization scheme for $-6 < X < -5$ and $13 < X < 14$. That these are much narrower than the conditional probability distributions shows the ‘overconfidence’ of the deterministic parametrization.




Figure 5. (a) Solid and dashed lines show the ranked probability skill scores for 75 initial condition ensemble forecasts at $t = 0.6$, based on differences between the Lorenz (1996) dynamical system and various parametrized versions of the system (see text for details), with $c = 10$ (solid) and $c = 4$ (dashed). The dotted line shows the ensemble mean RMS error for $c = 4$ (with values given on the right-hand side of the diagram). (b) Hellinger distance between the climatological probability distribution of the Lorenz (1996) dynamical system and the various parametrized versions, with $c = 10$ (solid) and $c = 4$ (dashed). Based on integrations over 400 time units.

Arnold *et al.* (2012). In the initial-value ensembles, evaluated at $t = 0.6$ time units (perhaps equivalent to about 3 days for weather forecasting), the initial conditions $X_k(t = 0)$ are known perfectly; hence there is no initial uncertainty, only simulator uncertainty. The solid line denotes the results with $c = 10$, the dashed line gives results with $c = 4$. For the initial-value problem, we use the ranked probability skill score (RPSS; Wilks, 2006) to assess the probabilistic skill in forecasting X_k . For the climate integrations, we use the Hellinger distance (related to the more familiar Kolmogorov–Smirnov distance; Pollard, 2002) between the ‘true’ and simulated probability distribution of X_k values. Note that the larger the RPSS, the more skilful is the forecast, whereas the smaller the Hellinger distance, the closer is the simulated probability distribution to the probability distribution of truth. Again, see Arnold *et al.* (2012) for details. Additionally, for the initial-value ensembles (Figure 5(a) for $c = 4$) we also show the traditional deterministic score, root mean square (RMS) error, averaged over all the individual forecasts.

A number of interesting results can be concluded from Figure 5:

1. Based on RPSS and Hellinger distance, and as expected, the $c = 10$ system is ‘easier’ to parametrize than the $c = 4$ system, and while stochastic parametrization improves forecast skill for both values of c , the improvement is relatively small when $c = 10$. By analogy, we would expect comprehensive weather simulators to be harder to parametrize deterministically, if their truncation scales probe the $-5/3$ part of the spectrum. As discussed above, there

is an inherent tension (perhaps one would even say incompatibility) between high-resolution simulation and deterministic parametrization.

2. Based on RPSS and Hellinger distance, there is an overall strong correlation between simulator performance in initial value mode and in climate mode, consistent with the philosophy underpinning the notion of seamless prediction. That is to say, the performance of the simulator in climate mode can be gauged by its success in initial-value mode. Of course, in the real world, one would not expect a one-to-one correspondence between weather and climate skill, because there are many slow climate processes which are not important for weather prediction. Nevertheless, the results here hint that skill on the weather time-scale should be considered a necessary step for reliable climate prediction.
3. The link between initial-value skill and climate accuracy is only apparent when probabilistic measures of skill are used to assess the initial value ensembles. If the more traditional deterministic RMS error metric is used to assess initial value skill, there is no correlation between initial value skill and climate skill; indeed the simulator with deterministic parametrization appears most ‘skilful’. As discussed in section 6, the conclusion to draw from this result is not that the link between weather and climate skill is metric dependent, but rather that the RMS error may actually be an inappropriate metric of weather forecast skill. The physical reason for this is discussed in section 6, where it is concluded that assessing simulators based on weather forecast RMS error may in fact be detrimental to the development of reliable climate forecast systems.
4. Based on RPSS and Hellinger distance, there is an overall advantage for the red-noise parametrization over the white-noise parametrization. This is consistent with the discussion above: in stochastic parametrization, it is necessary to represent the means by which uncertainty in the representation of subgrid processes affects the large-scale flow, on spatial scales larger than the simulator’s truncation scale, and on time-scales longer than the simulator’s time step. In Lorenz (1996), correlations between neighbouring X_k variables are small, and, for this particular model, there is not much benefit to the introduction of ‘spatially correlated’ noise. However, as Figure 5 shows, there is benefit in representing ‘temporally correlated’ noise. In general, for weather and climate simulators, one would expect the noise to be both spatially and temporally correlated.
5. There is an overall advantage for the multiplicative noise parametrization. This multiplicative noise parametrization is essentially that developed and tested in the ECMWF simulator by Buizza *et al.* (1999).

In the latest version of the ECMWF multiplicative noise scheme (or SPPT: stochastically perturbed parametrization tendency scheme; see Palmer *et al.*, 2009), the parametrization is given by

$$\dot{X}^{\text{stoch}} = (i + r^{\text{spec}} \mu) \dot{X}^{\text{det}}, \quad (4)$$




Figure 6. Realizations of the stochastic pattern generator used in the ECMWF stochastically perturbed parametrization tendency scheme (Palmer *et al.*, 2009). Strong (faint) lines correspond to positive (negative) values.

where \dot{X}^{stoch} denotes the stochastic tendency, \dot{X}^{det} the total deterministic tendency, and r^{spec} denotes a stochastic spectral pattern generator based on an uncorrelated series of red-noise processes, one for each spherical harmonic coefficient. The relative amplitude of these red-noise processes in spectral space is such as to produce Gaussian correlations in physical space (see Figure 6). In the results discussed below, there are two sets of such red-noise processes: one with 6 h decorrelation time, the other with smaller amplitude and 30-day decorrelation time (see Palmer *et al.*, 2009, for details). Finally, μ is an *ad hoc* parameter which clips the stochastic tendencies in the stratosphere and in the boundary layer. We return to this ‘ μ ’ parameter later.

A more overt example of the need to consider the representation of subgrid uncertainty on the resolved spatial scales arises in the stochastic backscatter scheme (Shutts, 2005; Berner *et al.*, 2009):

$$F_{\psi} = \left(\frac{b_{\text{R}} D_{\text{tot}}}{B_{\text{tot}}} \right)^{1/2} P_{\psi}. \quad (5)$$

Here the stream function forcing F_{ψ} is associated with an upscale energy transfer when, for example, divergent kinetic energy associated with deep convection is converted to rotational kinetic energy during mesoscale organization. This forcing is represented by a stochastic pattern generator P_{ψ} (either the spectral generator, cf. Figure 6, or an alternative cellular automaton – it can be noted in passing that cellular automata provide computationally cheap means to communicate information at the subgrid level, between adjacent grid boxes). Here D_{tot} denotes the diagnosed energy dissipation from the corresponding deterministic parametrizations, and B_{tot} and b_{R} are parameters which ensure dimensional consistency and degree of energy backscatter respectively.

Figure 7 (from Palmer *et al.*, 2009) shows the impact of SPPT on the probabilistic skill of medium-range forecasts of 850 hPa temperature in the Tropics using the ECMWF Ensemble Prediction System (EPS). The results are dramatic. The skill at day 2 of the probabilistic forecasts without stochastic parametrization is reached at day 6 with stochastic parametrization. It is hard to imagine any parametrization having such an effect on forecast skill.

The introduction of stochastic parametrization into the ECMWF simulator has fundamentally changed the skill of the EPS in more ways than one. Importantly, it has allowed




Figure 7. Continuous ranked probability skill score for 850 hPa temperature in the Tropics based on the ECMWF Ensemble Prediction System with no representation of model uncertainty (dotted line); the original ‘stochastic physics’ scheme of Buizza *et al.* (1999) (dashed line); the 2-time Stochastically Perturbed Parametrization Tendency scheme described in Palmer *et al.* (2009) (solid line). See Palmer *et al.* (2009) for details.

the estimation of initial uncertainty to be made using ensembles of (4D Var) data assimilations (EDA; Isaksen *et al.*, 2010). Until recently, EPS initial perturbations were made exclusively using singular vector analysis (e.g. Buizza and Palmer, 1995). The reason for this was that if an EPS was based solely on initial perturbations from ensembles of analyses, these perturbations had to be artificially inflated in order that EPS spread and skill matched in the medium range. Introduction of stochastic parametrization into the data assimilation process (and the use of higher resolution and hence less damped simulators) has enabled ensemble data assimilation to be used to generate initial EPS perturbations. Indeed Figure 8 shows the performance of EDA in terms of the relationship between ensemble spread at $T + 6\text{h}$ and ensemble mean error. It can be seen that, with representation of observation error only, not only is the EDA underdispersive but also the EDA spread does not discriminate well between low-error and high-error short-range forecasts (it is particularly underdispersive for high-error forecasts). By contrast, including SPPT and backscatter into EDA, not only is the overall level of spread much closer to that of error but the EDA spread now discriminates well between low- and high-error short-range forecasts.




Figure 8. Relationship between $T + 6$ h ensemble spread and $T + 6$ h ensemble-mean error of ensembles of data assimilations of the ECMWF forecast system, for tropospheric vorticity in the Northern Hemisphere, binned on error and based on: (dashed line) an ensemble with representation of observation uncertainty but not model uncertainty; (solid line) an ensemble with representation of observation and model uncertainty (based on stochastically perturbed tendencies and stochastic kinetic energy backscatter). The dotted line shows the ideal relationship. From Bonavita (2011).

Figure 9 shows that EPS-based probabilistic predictions of rainfall over Europe in the medium range are now extremely reliable.

There is no doubt that ensemble forecasts with stochastic parametrization are skilful, but are they more skilful than forecasts using the more traditional multi-simulator concept? This question, applied to the climate prediction problem, lies at the heart of this paper. Table 1 shows a comparison of probabilistic skill on the monthly time-scale (where copious verification data exist), based on three ensemble forecast systems (see Weisheimer *et al.*, 2011, for details). The first system is a multi-simulator ensemble comprising the climate simulators that contributed to the ENSEMBLES multi-simulator ensemble (Weisheimer *et al.*, 2009). The second system is the single-simulator ECMWF seasonal ensemble forecast system with stochastic (SPPT and backscatter) parametrization. The third ensemble is again based on the single-simulator ECMWF seasonal ensemble forecast system as above, but with no representation of simulator uncertainty (i.e. only initial uncertainty).

Results show that for seven of the eight binary forecast events considered (based on climatological temperature and precipitation terciles over all land points), the single-simulator ensemble with stochastic parametrization outperforms the multi-simulator ensemble. For one of the eight events, the skill estimates for the stochastic parametrization

ensemble and the multi-simulator ensemble only differ by the third significant digit. It might be imagined that the key reason that the single-simulator stochastic parametrization ensemble outperforms the multi-simulator ensemble is that the former has been made with a world-leading weather simulator. However, if we compare the skill of the multi-simulator ensemble with the skill of the same single-simulator ensemble without any representation of simulator uncertainty, then it can be seen from Table 1 that the latter is much the least skilful of the three ensembles for all events considered. This indicates that the single-simulator stochastic parametrization ensemble is not more skilful than the multi-simulator ensemble because this particular simulator is somehow inherently better (e.g. in terms of its deterministic forecast skill) than the other simulators.

In Weisheimer *et al.* (2011) it was also shown that on longer seasonal time-scales stochastic parametrization still has the edge against the multi-simulator ensemble for precipitation forecasts, but not for forecasts of surface temperature. This suggests (see section 4 below) that development of the stochastic approach for the land surface and for the oceans is also likely to be required in the future. The skill of a perturbed-parameter ensemble was also tested




Figure 9. Reliability diagram for the ECMWF Ensemble Prediction System at $t = 4$ days for prediction of rainfall exceeding 1 mm per day over the European domain, based on verification from March to May 2011. The values against the dots give the number of occasions where probability forecasts within a given 10% range (and at 0% exactly) were made.

Table 1. Brier skill scores for probabilistic predictions for all global land area 2 m temperature and precipitation grid points, based on exceeding upper climatological tercile (warm/wet) and not exceeding lower tercile (cold/dry) events for: the ENSEMBLES multi-simulator ensemble (MSE), an ensemble using the ECMWF simulator with stochastic parametrization (SPE) and an ensemble using the ECMWF simulator without any representation of simulator uncertainty (CTRL). Bold figures indicate the system with the highest score. From Weisheimer *et al.* (2011).

	T2m				Precipitation			
	May		Nov.		May		Nov.	
	Cold	Warm	Cold	Warm	Dry	Wet	Dry	Wet
MSE	0.178	0.195	0.141	0.159	0.085	0.079	0.080	0.099
SPE	0.194	0.192	0.149	0.172	0.104	0.118	0.095	0.114
CTRL	0.147	0.148	0.126	0.148	0.044	0.061	0.058	0.075

by Weisheimer *et al.* (2011). The skill scores turned out to be poor, but one cannot rule out the possibility that this was because the simulator in which the parameters were perturbed was not state-of-the-art for monthly and seasonal prediction. Further tests are needed within, for example, the ECMWF system, to evaluate the perturbed-parameter method. It is certainly not inconceivable that some combination of perturbed-parameter and stochastic parametrization techniques may prove optimal.

A key property of stochastic parametrization is its potential ability to influence the mean state of the simulator and hence reduce the mean bias of the simulator against observations. That is to say, the interaction of the imposed noise with the nonlinearity of the simulator can generate a ‘rectified’ time-mean response. In this way, it is possible that stochastic parametrization can help alleviate some of the systematic biases of climate simulators. Figure 10 shows an example of such alleviation (from Berner *et al.*, 2012, who also show a positive impact of stochastic backscatter on the mean state of simulations in the Tropics).

However, a problem revealed by Figure 10 is that the impact of stochastic parametrization on simulations of Northern Hemisphere circulation is very similar to the impact of either increasing simulator resolution (i.e. modifying the dynamical core), or modifying the conventional deterministic parametrization schemes. Dynamical reasons for this ‘degeneracy’ are discussed in Palmer and Weisheimer

(2011). These explain why improving the fidelity of climate simulators has been so difficult over the years, and why it is very easy for a simulator code to contain many sets of ‘compensating errors’. This is a key reason why data assimilation can provide such a powerful tool for enabling simulator development while minimizing such compensating-error problems (see Palmer and Weisheimer, 2011, for a discussion). This problem of degeneracy is discussed further in section 4 below.

4. Stochastic parametrization at the process level

Despite these rather positive results, stochastic parametrization is still at a rudimentary state of development: the stochastic parametrization concept described above has only been applied to the atmospheric component of coupled simulators. There is clearly a need to extend the concept to the oceans, the land surface, the cryosphere, the biosphere and so on. The techniques which can be used to develop stochastic parametrizations are manifold, and the logic inductive rather than deductive. A technique of particular relevance is the type of coarse-grain analysis developed in Frederiksen and Kepert (2006) and Shutts and Palmer (2007). Moreover, the sort of experimental programmes advocated by Jakob (2010) are just as important for the development of stochastic parametrization as for deterministic.




Figure 10. Mean systematic error of 500 hPa geopotential height fields for extended boreal winters (December–March) of the period 1990–2005. Errors are defined with regard to the observed mean field (contours), consisting of a combination of ERA-40 (1990–2001) and operational ECMWF analyses (2002–2005). Shown are the systematic error of experiments: (a) low-resolution T95 simulator, (b) T95 simulator with stochastic kinetic-energy backscatter, (c) high-resolution T511 simulator and (d) T95 simulator with improved deterministic parameterizations. Contour interval 2 dm. From Berner *et al.* (2012). © American Meteorological Society. Used with permission.

However, even for the atmospheric component of climate simulators, there is a need for uncertainty to be incorporated in the development of parametrization at the process level, rather than as a 'bolt-on extra'. For example, in describing the multiplicative noise parametrization in section 3, reference was made to the *ad hoc* parameter μ which clipped the stochastic noise both in the boundary layer and in the stratosphere. The parameter was introduced for plausible reasons, but also because it improved forecast scores. However, one should not introduce parameters purely because of empirical pragmatism: they must additionally have some basis in science. For the stratosphere, the scientific basis is not hard to find. Much of the diabatic heating in the stratosphere is associated with infra-red radiation emitted by carbon dioxide molecules. However, unlike water, carbon dioxide is well mixed in the atmosphere and there is little subgrid variability. Hence there is no need to represent this process stochastically. It is also conceivable that, at least in sufficiently homogeneous terrain well away from orography, a typical boundary layer 'eddy' associated with surface form drag is also sufficiently small in scale that grid-scale stochasticity in grid-scale vertical mixing will be relatively small. This argues that, instead of having an *ad hoc* μ parameter, aspects of stochastic parametrization should be developed at the process level.

The case for stochastic parametrization at the process level is fairly clear when discussing processes like convection (e.g. Lin and Neelin, 2003; Plant and Craig, 2008), and imaginative new stochastic schemes for parametrizing different convective cloud families are being developed using cellular automata (e.g. Bengtsson-Sedlar *et al.*, 2011) or stochastic lattice models (Khouider *et al.*, 2003; Frenkel *et al.*, 2011). However, even something as basic (and in principle well known) as radiation needs stochastic treatment; gridbox surface radiative fluxes can depend strongly on poorly resolved near-grid-scale circulations. For example, under a region of stratocumulus, surface fluxes will depend strongly on whether in-cloud shallow convection is of the closed cell or open cell type. It is unrealistic to expect these small-scale circulations to be deterministic functions of the large-scale variables; such effects therefore represent a source of uncertainty in forecasts of surface temperature that should be incorporated at the process level into the simulator equations.

However, there is a separate and quite fundamental argument for the need to develop stochastic parametrization as an inherent part of simulator development, and not as a 'bolt-on' extra. In section 3 it was shown that stochastic parametrization had an impact on a simulator's systematic error. Consider the implications of this for setting the parameters α of the deterministic parametrizations $P(X_k, \alpha)$.

For example, the parameter often called 'convective entrainment' represents the strength of the process whereby environmental air is entrained laterally into convective plumes. It is well known that climate simulations can be especially sensitive to the value of this parameter (Stainforth *et al.*, 2005). However, if the notion of a subgrid ensemble of convective plumes is not well founded due to power-law structure and associated scale invariance, then neither is the existence of a well-defined value for the convective entrainment parameter. As such, and this is universally recognized by the scientists who develop climate simulators, the values of these parameters have to some extent to be 'tuned' based on the fit of simulator output to sets of

observations of the large-scale structure of the atmosphere (either based on weather forecasts or climate integrations).

However, consider the implications of such tuning exercises if 'bolt-on' stochastic parametrizations change the mean state of the simulator. They imply that values of the parameters α_{det} which have been optimally tuned for a deterministic simulator will not be optimal in a stochastic simulator. This implies that α_{det} are not in fact optimal at all. (See also Tompkins and Berner, 2008.)

This situation is familiar in many other areas of physics. Consider the vertical motion of a table-tennis ball with mass m_0 inside water. As found by Green in the 19th century, the motion of the ball obeys Newton's law $F = ma$, where F is the Archimedean buoyancy force, but where $m = m_0 + M/2$ and M is the mass of water occupying the same volume as the table-tennis ball. In other words, while in the absence of randomly fluctuating molecules of water the motion of the ball obeys $F = m_0a$, in the presence of these random fluctuations the motion of the table-tennis ball behaves as if it were half full of water. This 'renormalization' of mass has measurable effects: the initial acceleration of the ball back towards the surface is about seven times smaller than it would otherwise be. Quantum field theorists recognize in these arguments the difference between the 'bare' and 'effective' mass of a particle such as an electron in the presence of the fluctuating photonic field.

In the same way, we argue here (as indeed was argued by Frederiksen and Davies, 1997; Frederiksen, 1999) that parameter tuning for weather and climate simulators must be done in the presence of parametrized representations of the inherent stochasticity associated with the scale-invariant properties of the underlying equations. The notion of 'bolt-on' stochastic parametrization (for use when the simulator is run in probabilistic ensemble mode) using deterministically pre-tuned parameter values is not a scientifically sound procedure.

The focus of attention in this paper has largely been on the parametrization of physical processes. It has been argued that even when we know the underpinning equations with accuracy the resulting parametrizations should be considered stochastic. However, for parametrizing other processes (e.g. biological, chemical and perhaps aerosol), where the underpinning equations are not known with accuracy, the need for stochastic representations is no less important and necessary. At the very least, if different deterministic closures $\{A_p, B_p, \dots, C_p\}$ have been proposed for process P , and observations cannot rule out any one of these closures over another, a particular closure could be chosen randomly at a given grid box and time step, using the types of spatially and temporally correlated pattern generators discussed above.

It is important to emphasize that none of the above implies that we must not continue to develop, refine, improve and extend our subgrid parametrizations. This work remains as critical in the future as it has been in the past. However, it is argued here that this development, refinement, improvement and extension should be performed within an inherently probabilistic, and hence stochastic, parametrization framework. The author believes that research and development within this more general framework will allow innovative ideas to flourish and parametrization breakthroughs to occur.

5. Dynamical cores and stochastic processors

It was noted above that stochastic parametrization inevitably involves the representation of the upscale propagation of subgrid uncertainty onto the resolved grid. This suggests that, just as it may be futile trying to develop precise deterministic parametrizations, so also it is futile to develop precise deterministic dynamical cores, especially for the evolution of scales near the truncation scale.

While this may be the case, if we have accounted statistically for upscale propagation of uncertainty in the parametrization, then the only disadvantage to retaining a precise deterministic dynamical core is the computational burden. However, how would one go about defining a probabilistic dynamical core which is both consistent with the equations of motion, and would provide a significant reduction in computational cost compared with current deterministic cores? After all, computing a stochastic field which comprises large numbers of pseudo random-number generators is certainly not computationally cheap.

However, there is an emerging technology that may present a way forward here and at the same time provide a new type of synergy between software development (of the high-resolution probabilistic Earth-system simulator) and the very hardware needed to integrate a simulator's equations. This technology (e.g. Palem, 2005) is motivated by the fact that a significant fraction of a conventional computer's energy consumption is associated with heat dissipation at the chip level. Hence, if the processors of a computer could be designed so that when the voltage across the individual transistors is reduced, the computer would operate with significantly reduced energy consumption but at marginally reduced (e.g. 99% instead of 100%) accuracy, this capability would certainly be worth exploiting.

Indeed the issue that bit-reproducible computation may become a thing of the past is beginning to be recognized in the supercomputing industry too. In a recent presentation on challenges in application scaling in an exascale environment, IBM's Chief Engineer noted (http://www.ecmwf.int/newsevents/meetings/workshops/2010/high_performance_computing_14th/index.html) that increasingly there will be 'a tension between energy efficiency and error detection', and asked whether there needs to be a new software construct which identifies critical sections of code where the right answer must be produced – implying that outside these critical sections errors can (in some probabilistic sense) be tolerated.

One can perhaps imagine a future energy-efficient computer with clusters of processors each with different levels of accuracy, integrating a future-generation dynamical core. The more accurate the processor, the larger will be the scales of motion for which it computes tendencies. The 'right answers' will be produced only for the large-scale tendencies. Since, overall, computations are dominated by estimation of tendencies nearer the truncation scale, the synergistically designed probabilistic supercomputer need have relatively few of these slower energy-intensive processors. On top of this, the stochastic parametrizations themselves would be computed using the energy-efficient probabilistic chips.

There is a link here to the work of Lander and Hoskins (1997), who argued that sophisticated and computationally expensive parametrization schemes should only be applied to the more 'believable' scales in a simulator, i.e. scales far removed from the truncation scale. They propose that

simpler parametrization schemes could be used on the 'unbelievable' scales near the truncation scale. This idea has some resonance with the proposal discussed here whereby less-believable computations near the truncation scale could be executed on relatively fast energy-efficient probabilistic processors, leaving computations at the large 'believable' scales for traditional energy-intensive bit-reproducible processors.

What would prevent utterly erroneous computations from compromising the validity of the computation (e.g. due to big errors in the exponents of key real numbers representing the small scales)? It could be prevented partly by performing numerical checks against prior physical bounds, and partly by repeating computations several times, and taking the mode of some small ensemble of obtained values. This has to be a matter for future research, providing hardware developments look sufficiently promising.

As well as being more energy efficient, it is possible that such probabilistic architectures may offer a significant increase in computational speed-up for climate simulator codes. If this is so, probabilistic processing may allow a route to cloud-resolved climate simulation much faster than anyone had previously expected, again allowing one to realize the goals of the World Summit on Climate Modelling (Shukla *et al.*, 2010) in the foreseeable future.

The use of probabilistic Earth-system simulators running on machines built with stochastic processors, i.e. where the inherent quantum-mechanical noise associated with electrons flowing through transistors becomes a resource rather than a nuisance, provides a new synergy between software and hardware design in the field of weather and climate prediction, hitherto unimagined.

6. Probabilistic forecasting and seamless weather prediction: opportunities and possible obstacles

Following Bjercknes (1904), NWP has historically been considered an example of a deterministic initial-value problem. The notion of probabilistic forecasting using ensemble prediction methods has evolved more recently as a tool to mitigate the effects of chaotic weather variability. Operational ensemble weather prediction systems have been implemented since the mid 1980s (Murphy and Palmer, 1986), long before ensemble methods became commonplace in climate prediction (see the review by Lewis, 2005). Following considerable investment in ensemble prediction at a number of NWP centres since these early days, most NWP centres now develop both a high-resolution deterministic forecast system and a lower-resolution EPS, and strategic goals are targeted separately on improvements in both deterministic scores for the high-resolution system (e.g. RMS error or anomaly correlation coefficient) and probabilistic scores for the EPS (e.g. ranked probability skill score).

These goals are individually challenging and require the determined effort of scientists across a range of disciplines (numerics, parametrization, data assimilation, etc). This presents important questions about allocation of resources. How, for example, should an NWP centre partition its human resources to meet both the deterministic strategic goal on the one hand and the probabilistic target on the other? A common view is that if most human resources are put into meeting the deterministic goal, the resulting improvements to the deterministic forecast system will




Figure 11. 500 hPa geopotential anomaly correlation coefficient over Northern Hemisphere Extratropics for March–May 2011. Light solid line: high-resolution (T1279) ECMWF deterministic forecast. Dashed line, unperturbed control forecast from the (T639) ECMWF ensemble prediction system; dotted line, deterministic forecast based on the ensemble average over the members of the ECMWF ensemble prediction system.

necessarily benefit the EPS and help ensure the probabilistic target is also met.

Unfortunately, this concept of ‘trickle-down’ does not apply to the development of stochastic parametrization. Weather simulators with stochastic parametrizations cannot produce forecasts with as low RMS error, or as high anomaly correlation coefficient, as equivalent simulators with deterministic parametrizations (see Figure 5(a) for an explicit illustration of this in the parametrized Lorenz ‘96 model). The reason why a probabilistic simulator will not outperform a comparable deterministic simulator in terms of deterministic scores is similar to the reason why the most skilful ‘deterministic’ forecasts are associated with the ensemble-mean forecast (see Figure 11). The reason why an ensemble-mean forecast has especially high deterministic skill is that the relatively unpredictable components of the flow are ‘damped out’ in an ensemble-mean field. However, a penalty is paid for such dynamical smoothing. An ensemble-mean forecast is unlikely to predict the occurrence of a severe weather event, if such an event is relatively unpredictable; the ensemble-mean forecast ‘hedges’ towards climatology and away from such events. Now, as discussed in sections 2 and 3, a deterministic bulk-formula parametrization can be considered as providing an estimated mean tendency based on a putative ensemble of inherently unpredictable subgrid processes, and hence will produce a ‘damped’ simulation of the flow at sub-synoptic scales. In the same sense that an ensemble-mean forecast has low deterministic error, a simulator with deterministic bulk-formula parametrization will tend to produce forecasts with lower RMS error than an equivalent simulator with stochastic parametrization, particularly for near grid-scale circulations; recall (cf. Figure 3) that each realization of the stochastic parametrization is designed to represent a potential realization of the subgrid flow, rather than an ensemble average. However, as with the ensemble-mean forecast, there is a price to pay for this smoothing: a tendency for the simulator to hedge away from simulating extreme flows. This effect will obviously be strongest for small scales. However, as discussed in section 2.2, small-scale errors can

be expected to propagate, intermittently but rapidly, to larger-scale components of the flow.

Put bluntly, stochastic parametrization is anathema to the strategic goal of maximizing deterministic skill! As such, development of stochastic parametrization at the process level, the type of activity discussed in section 4, will not naturally emerge from research that is focused primarily at improving the high-resolution deterministic forecast system.

Should NWP centres therefore start planning for the day where they focus exclusively on developing probabilistic forecast systems, drop their higher-resolution deterministic predictions, and measure progress primarily in terms of improvements to probabilistic scores?

Some may argue against this, noting that the enhanced skill of higher-resolution deterministic forecast systems justifies their continued separate development. Unquestionably, this was true in the past: in the early days of operational EPS, the deterministic skill of the unperturbed EPS control forecast was substantially poorer than that of the higher-resolution deterministic forecast. However, these days, as shown in Figure 11, the skill of the higher-resolution deterministic forecast is no longer substantially greater than that of the EPS. It should certainly not be concluded from this that there is no need for the development of high-resolution simulators. There is evidence that at T1279 resolution extreme weather events (such as hurricanes) can be simulated with greater realism than at T639 resolution. Rather, the point of Figure 11 is that the impact of high resolution is more subtle than it was in the past, and much less apparent in headline strategic scores such as the 500 hPa anomaly correlation coefficient. In section 2.2, the question was raised: how can we ensure that the advantages of integrating simulators at higher and higher resolution will not somehow be compromised by the intermittent upscale propagation of error? That is to say, how can we produce high-resolution forecast systems that are reliable (the overall theme of this paper)? The author believes that the answer to this question is that future high-resolution forecast systems must be explicitly probabilistic.

Others, arguing against this conclusion, may claim that weather forecast offices will continue to require high-resolution deterministic forecasts for the foreseeable future, since weather forecast customers demand precise deterministic forecasts, and find probabilistic forecasts difficult to understand and difficult to use. This argument becomes yet stronger when one realizes that the computational cost of a single high-resolution deterministic forecast is small compared with the cost of a full EPS.

Why is it that the public wants and expects detailed deterministic forecasts? Certainly, nobody wants an uncertain forecast if a perfect deterministic forecast is available. But the latter is not available and never will be. In the author's opinion, a key reason why the public expects deterministic forecasts is simply because that is what they have been given and hence led to expect, ever since the days of Fitzroy when the first weather forecasts were made available to the general public. However, in cases where uncertainty is routinely expressed to the public, e.g. in the US National Hurricane Center's 'cone of uncertainty' for hurricane track predictions (<http://www.nhc.noaa.gov/aboutcone.shtml>), the author's own informal research suggests that the public understands and indeed respects these uncertain predictions, and consequently no longer demands deterministic predictions.

Independently of whether the public are ready to accept the notion of an explicitly uncertain forecast, perhaps there is an argument that by focusing on probabilistic forecasting methods, the traditional skills of the weather forecaster will somehow be undermined. However, the author believes that the skills of human forecasters will be needed as never before when forecasts are primarily probabilistic in nature. In particular, there will be a need for a greater dialogue between forecasters and customers to help guide individual customers formulate weather-sensitive decision strategies appropriate to their circumstances. A simple (and rather idealized) example is based on the cost/loss model (Murphy, 1969). If a customer incurs a loss L if a particular weather event E (e.g. based on temperature, precipitation, wind, or some combination thereof) occurs, but can take protective action at cost C , then it makes rational sense to take this protective action on those occasions when the forecast probability of E exceeds C/L . In these circumstances, the job of the forecaster will be to 'tease out', at least approximately, the customer's C/L and therefore enable that customer to decide on the optimal threshold probability above which preventative action should be taken. Using this cost/loss model, Figure 12 shows the 'potential economic value' of the EPS, compared with that of the high-resolution deterministic forecast – the latter can be considered as a probabilistic forecast producing only probabilities of one or zero – based on precipitation events at forecast day 4. A 'potential economic value' of unity would correspond to a hypothetical perfect deterministic forecast, and a 'potential economic value' of zero would correspond to the value obtained by knowing only the climatological probability of E . The value of the EPS is substantially higher than that of the high-resolution deterministic forecast – indeed, for a range of users, the high-resolution deterministic forecast, by itself, has no value at all over and above a decision based only on the climatological probability of E . Once again, it should be stressed that this does not at all imply that there is no merit in high resolution. Rather, Figure 12 suggests that the value of high resolution is masked when assessed




Figure 12. Potential economic value (Murphy, 1969) as a function of user cost/loss ratio, based on prediction of rainfall exceeding 1 mm per day over the European domain for March–May 2011 (1 = value of a perfect deterministic forecast, 0 = value associated with a climatological probability forecast). Solid line for ECMWF Ensemble Prediction System; dotted line associated with ECMWF high-resolution deterministic forecast.

in deterministic mode. There is no reason to doubt that a T1279 EPS would have higher 'potential economic value' than the current T639 EPS, especially for severe events E .

In practice, decision strategies will be much more complex than suggested by a simple cost/loss models, for example requiring knowledge of the customer's 'utility function' which maps, usually nonlinearly, multiple correlated weather variables to some quantity relevant to the customer (number of ice creams sold, megawatts of electricity produced). It will be the job of tomorrow's weather forecaster to help the weather-sensitive customer to formulate his or her decision strategy in these realistic circumstances. It is interesting to note that, in this respect, great advances have been made recently in applying ensemble-based probability forecasts to provide flood risk assessments for farmers and community leaders in developing countries in the Tropics, and these have been shown to have genuine value in saving lives and property (Webster *et al.*, 2010).

In their interface with the general public, media forecasters need not only to be open about the inherent uncertainty in forecasts, but they should routinely relay to the public the fact that techniques exist to quantify this uncertainty. This does not necessarily mean displaying isopleths of probability on TV. However, during media forecasts, forecasters could refer to web sites, even better to interactive displays ('press your red button'), where 'fan charts' for temperature and rainfall, similar to those used by the Bank of England in forecasting inflation rate and gross domestic product (<http://www.bankofengland.co.uk/publications/inflationreport/irfanfch.htm>), can be displayed for key cities.

In conclusion then, it is proposed that, in the coming decade, NWP centres should start to focus exclusively on developing probabilistic forecast systems, dropping their separate higher-resolution deterministic forecast systems,

and, importantly, measuring progress, and formulating strategic goals, principally in terms of improvements to probabilistic scores. Such a strategy would certainly be consistent with the aims of forecast offices to provide reliable predictions of severe weather: no forecast can be considered reliable without an accurate assessment of forecast uncertainty, and severe weather events are often the most unpredictable and hence uncertain. The benefits of this, in addition to that of improving the reliability of weather forecasting *per se*, would be that improvements made to simulators on the weather time-scale would likely also improve the reliability of simulators for longer-term climate prediction.

7. Towards a seamless probabilistic Earth-system simulator for weather and climate prediction

As discussed in the Introduction, output from comprehensive climate simulators informs mitigation policies, climate adaptation strategies, efforts to understand the impacts of climate geoengineering, and generally reduces society's vulnerability to current and future climate. One is hard pressed to think of examples where computer code has such societal relevance! And yet, as discussed above, there are substantial challenges (theoretical, computational and human) that need to be overcome if we are to progress significantly to the goal of providing society with reliable estimates of future climate – regional and global.

In discussing possible ways to meet these challenges, consider by analogy the state of the European civil aircraft industry in Europe in the mid 20th century. At this time, all the major European countries produced their own civil aircraft. However, it was realized that aircraft were becoming too complex and too expensive for individual countries to develop and manufacture independently. Within this milieu, the Airbus consortium (<http://www.airbus.com/>) was formed. At the time, there must have been much agonizing at the national level as to whether national aerospace industries were doing the right thing getting together in this way. In retrospect, there can be little doubt but that it was. And so, within the Airbus consortium, these same national aerospace industries now focus on specific aspects of the design and production of aircraft in their fight for market share with their great US rival, Boeing.

Hence, by analogy, we can imagine a multi-national Earth-system simulator supported by teams of scientists from national climate and academic institutes. Different teams would focus on different aspects of the simulators: dynamical cores, oceans, clouds, aerosols. etc., and on the design of experiments which integrate these aspects together. All should contribute to the analysis and diagnosis of results. To support this, computational resources would be available, not only for operational integrations, but also for plentiful research experimentation. Results from the small number of simulators worldwide might continue to be combined in a multi-simulator ensemble, but since each is now based on stochastic-dynamic closure, the resulting ensemble would be much less prone to the type of systemic failure that current generation multi-simulator ensembles are capable of. National weather services would still play a crucial role in development work, in conducting scientific experiments, and in communicating the results from the science to their governments and society alike.

Is this a possible framework for the development of future Earth-system simulators? To some extent it already is. For example, within Europe, many climate institutes use the same (NEMO; <http://www.nemo-ocean.eu/>) ocean simulators. Indeed, development of the EC-Earth simulator (Hazeleger *et al.*, 2010) provides a specific example of how international cooperation can be successful, having been developed from the ECMWF seasonal forecast simulator, ECMWF itself being an outstandingly successful example of international cooperation in the context of NWP.

Given the merits of pooling resources, why would we not want to go further down this route of rationalization? The key argument for not adopting the 'Airbus' model is that we need extensive simulator diversity in order to estimate prediction uncertainty. However, the stochastic science discussed in the previous sections (and this is why the discussion has been so extensive) suggests that an alternative approach to representing simulator uncertainty is beginning to emerge, and, on time-scales where verification data exists, this alternative approach can outperform that provided by conventional multi-simulator ensembles. That is, the argument for maintaining the status quo of extensive simulator diversity is being undermined by scientific developments.

It should be stressed that it is not being suggested here that stochastic parametrization implies that all we need is one 'World Weather and Climate Simulator'. Airbus has undoubtedly been successful, not only because it can draw from the pooled resources of European aerospace industries but also because it has a competitor from another geopolitical grouping. Similarly, one would imagine that if there was some rationalization of climate simulator development effort, which embraced the notion of stochastic parametrization as the primary means to estimate simulator uncertainty, then we would still have enough (quasi-)independent Earth-system simulators to foster competition and creativity. What is a desirable number of comprehensive Earth-system simulators? This obviously depends on an assessment of the minimum human and computational resources needed to develop and maintain an Earth-system simulator. However, the author would broadly concur with the findings of the World Summit on Climate Modelling (Shukla *et al.*, 2010), that development of 'a small number' based around major geopolitical groupings might be ideal.

In the course of this paper, evidence has been given as to how the development of explicitly probabilistic weather and climate simulators will lead to more reliable estimates of uncertainty. At the beginning of the paper, it was also suggested that these methods might be able to actually reduce uncertainty. In considering this possibility, let us focus here on what must surely be the most important, as well as the most uncertain, of all the feedbacks in the climate change problem: that associated with cloud. As is well known (Solomon *et al.*, 2007), even the sign of the cloud feedback is uncertain.

One of the problems in thinking about the notion of 'cloud feedback' is that a world without cloud, and hence without cloud feedback, would be utterly alien to us: clouds are absolutely intrinsic to the circulation patterns we observe around us. Not only are clouds determined by the temperature and humidity structure associated with these circulation patterns, but also clouds in turn are key to determining these circulation patterns, both locally and

remotely. For example, anomalous latent heat release in convective cloud systems over the Caribbean may be key to setting up a blocking anticyclone over Europe, while the stratus decks that form locally in the vicinity of the blocking anticyclone are key to determining the surface temperature under the block.

This means that we cannot treat the problem of cloud feedback solely as a problem in atmospheric thermodynamics; the problem is as much dynamic as thermodynamic. For the sake of argument, let us consider climate as a dynamical system with distinct nonlinear regime structures (Palmer, 1998; Straus *et al.*, 2007) in both the Tropics and Extratropics. These regimes will in turn have distinct cloud properties (Williams and Webb, 2009): a blocking anticyclone may be dominated by relatively thin stratus clouds in winter and cirrus clouds in summer, while a cyclonic weather regime will contain significant amounts of thick nimbostratus cloud at all times of year. From this dynamical perspective, a key element of understanding the cloud feedback problem lies in estimating reliably how anthropogenic forcing will change the frequencies of occurrence of the regimes. (Changes to the structure of the stability of the regimes, this may be a secondary aspect of the problem.) That is to say, changes in these frequencies of occurrence will be one of the key factors in determining whether upper- or lower-level clouds increase or decrease as a result of anthropogenic climate change. Small wonder, then, that current climate simulators have such difficulty in simulating the sign of cloud feedback with any consistency. As discussed above, these same simulators have difficulty simulating the statistics of observed weather regimes.

Hence, to really make progress in reducing the uncertainty in cloud feedbacks it will be essential that the statistics of weather regimes are simulated correctly: their three-dimensional structure, their embedded cloud properties and their frequency of occurrence (see also Stephens, 2005). This is a profoundly challenging dynamical problem, and results suggest that the current generation of climate simulators is not fully up to the challenge.

The same arguments could be applied to another of the important uncertainties in climate prediction: the impact of aerosols. Here the key uncertainties relate to the indirect effect of aerosols, i.e. through their modification of cloud. Again, this indirect effect will be regime dependent, implying that we will never be able to assess aerosol impact reliably in the atmosphere without an accurate simulation of structure and frequency of occurrence of weather regimes.

With this in mind, we can suggest why the proposal for inherently probabilistic Earth-system simulators will reduce uncertainty in predictions of climate:

- (a) As discussed above, representing simulator uncertainty by stochastic parametrization undermines the inherent need for a large diversity of simulators, meaning that it will be possible to pool human and computational resources. Economies of scale will enable climate scientists to have dedicated access to top-of-the-range supercomputers, enabling key physical processes to be simulated, including *in situ* Rossby wave breaking, key for maintaining weather regimes against dissipation (Woollings *et al.*, 2008), and remote tropical convective systems which help 'force' these regimes.
- (b) Being more consistent with the underlying equations of motion, it could be argued that if there are to be breakthroughs in parametrization, e.g. of the effects of unresolved cloud systems, they are more likely to occur within a more general probabilistic framework, than within the traditional deterministic framework.
- (c) Development of seamless probabilistic weather and climate simulators will enable sophisticated diagnostic tools from data assimilation to be used to reduce climate prediction uncertainty (Rodwell and Palmer, 2007), e.g. based on studies of biases in analysis increments, composited on specific weather regimes. The use of data assimilation in assessing stochastic parametrization was illustrated in Figure 8.
- (d) There is evidence that stochastic parametrizations can improve directly estimates of the frequency of occurrence of weather regimes (Jung *et al.*, 2005). The reason relates to the rectification of the flow by stochastic noise. As a simple analogy, imagine a ball bearing moving in a potential with multiple minima; an overly damped system will lead to the ball bearing spending too much time in the dominant well and this will be reflected in a bias in the time-averaged position of the ball.

8. Conclusions

Compared with the economists, weather and climate scientists do indeed know their equations, at least as they relate to the physics of weather and climate. However, these equations cannot be solved by pencil and paper. Algorithmic representations of the equations of motion necessarily involve errors, and with conventional numerical algorithms based on deterministic closures these errors appear to lead to substantial biases and considerable uncertainty in simulating climate. Some discussion has been given to the possibility that convergence to the 'true' underlying equations with increasing resolution may be exceptionally slow, due to the '-5/3' power law for atmospheric energy. Some technical discussion has been given to an alternative strategy for closing the equations, where the inherent uncertainty in any algorithmic representation of the underlying equations is recognized explicitly. It is suggested that breakthroughs in the parametrization problem, if they are to occur, will be more likely within a stochastic framework than in the traditional deterministic framework.

On time-scales where verification data exist, these stochastic methods are beginning to outperform conventional multi-simulator ensembles. However, there is much work to be done before all relevant Earth-system parametrizations can be said to have been developed in this probabilistic way. Indeed, it has been concluded that focusing excessively on the traditional challenge in NWP, of reducing deterministic measures of forecast error, may increasingly become an obstacle to the seamless development of reliable probabilistic weather and climate simulators. It was argued that it may indeed be time to consider focusing operational weather forecast development entirely on high-resolution ensemble prediction systems.

A key aspect of this paper has been discussion on some of the implications of a move towards probabilistic Earth-system simulation – implications that transcend the technical aspects of stochastic parametrization. In particular, by undermining the argument for a large pool of quasi-independent simulators, the stochastic parametrization

programme provides new support for one of the key conclusions of the World Summit on Climate Modelling (Shukla *et al.*, 2010): for a pooling of human and computational resources amongst climate institutes and for a substantial rationalization of development work towards a very small number of independent Earth-system simulators.

Given the importance and urgency of predicting Earth's climate as accurately as science and technology allow, it is time to give serious thought to such change.

Acknowledgement

My thanks to a number of colleagues who helped produce the results shown in this paper, and with whom I had valuable discussions and received helpful comments. These include: James Annan, Hannah Arnold, Peter Bauer, Judith Berner, Roberto Buizza, Tony Busalacchi, Erland Källén, Chris Farmer, Anna Ghelli, Brian Hoskins, Hugh McNamara, David Richardson, Gregory Seregin, David Straus, Andrew Majda, J. Shukla, Julia Slingo, Rowan Sutton, Adrian Tompkins, Jean-Noël Thépaut, Alan Thorpe and Antje Weisheimer. I am also grateful to two anonymous reviewers for their comments. My thanks also to Rob Hine for helping produce the illustrations.

References

- Arnold H, Moroz I, Palmer TN. 2012. Stochastic parametrizations and model uncertainty in the Lorenz '96 system. *Phil. Trans. R. Soc.* (submitted).
- Arrhenius S. 1896. On the influence of carbonic acid in the air on the temperature of the ground. *The London, Edinburgh and Dublin Philosophical Magazine and Journal of Science Series 41*: (reprinted in *The Warming Papers*, 2011, Archer D, Pierrehumbert R (eds). Wiley-Blackwell: Chichester).
- Bengtsson-Sedlar L, Körnich H, Källén E, Svensson G. 2011. Large-scale dynamical response to sub-grid scale organization provided by cellular automata. *J. Atmos. Sci.* **68**: 3132–3144.
- Berner J, Shutts GJ, Leutbecher M, Palmer TN. 2009. A spectral stochastic kinetic energy backscatter scheme and its impact on flow dependent predictability in the ECMWF ensemble prediction system. *J. Atmos. Sci.* **66**: 603–626.
- Berner J, Jung T, Palmer TN. 2012. Systematic error model error: the impact of increased horizontal resolution versus improved stochastic and deterministic parameterizations. *J. Climate* (in press).
- Bjerknes V. 1904. Das Problem der Wettervorhersage betrachtet von Standpunkte der Mechanik und der Physik. *Meteorol. Z.* **21**: 1–7.
- Bonavita M. 2011. Ensemble of data assimilations and estimation of uncertainty. In *ECMWF 2011 Seminar Proceedings*.
- Buizza R, Palmer TN. 1995. The singular vector structure of the atmospheric global circulation. *J. Atmos. Sci.* **52**: 1434–1456.
- Buizza R, Miller M, Palmer TN. 1999. Stochastic representation of model uncertainties in the ECMWF ensemble prediction system. *Q. J. R. Meteorol. Soc.* **125**: 2887–2908.
- Doblas-Reyes FJ, Balmaseda MA, Weisheimer A, Palmer TN. 2011. Decadal climate prediction with the ECMWF coupled forecast system: impact of ocean observations. *J. Geophys. Res.* **116**: D19111, DOI: 10.1029/2010JD015394.
- Frederiksen JS. 1999. Subgrid-scale parameterizations of eddy-topographic force, eddy viscosity, and stochastic backscatter for flow over topography. *J. Atmos. Sci.* **56**: 1481–1494.
- Frederiksen JS, Kepert SM. 2006. Dynamical subgrid-scale parameterizations from direct numerical simulations. *J. Atmos. Sci.* **63**: 3006–3019.
- Frederiksen JS, Davies AG. 1997. Eddy viscosity and stochastic backscatter parameterizations on the sphere for atmospheric circulation models. *J. Atmos. Sci.* **54**: 2475–2492.
- Frenkel Y, Majda AJ, Khouider B. 2012. Using the stochastic multcloud model to improve tropical convective parametrization: a paradigm example. *J. Atmos. Sci.* **69**: 1080–1105.
- Goldstein M, Rougier JC. 2004. Probabilistic formulations for transferring inferences from mathematical models to physical systems. *SIAM J. Sci. Comput.* **26**: 467–487.
- Hansen J, Russell G, Rind D, Stone P, Lacis A, Lebedeff S, Ruedy R, Travis L. 1983. Efficient three-dimensional global models for climate studies. *Mon. Wea. Rev.* **111**: 609–662.
- Hasselmann K. 1976. Stochastic climate models, Part I: Theory. *Tellus* **28**: 473–485.
- Hazeleger W, Severijns C, Semmler T, Ştefănescu S, Yang S, Wang X, Wyser K, Dutra E, Baldasano JM, Bintanja R, Bougeault P, Caballero R, Ekman AML, Christensen JH, van den Hurk B, Jimenez P, Jones C, Källberg P, Koenigk T, McGrath R, Miranda P, Van Noije T, Palmer T, Parodi JA, Schmith T, Selten F, Storelvmo T, Sterl A, Tapamo H, Vancoppenolle M, Viterbo P, Willén U. 2010. EC-Earth: a seamless Earth-system prediction approach in action. *Bull. Am. Meteorol. Soc.* **91**: 1357–1363.
- Isaksen I, Bonavita M, Buizza R, Fisher M, Haseler J, Leutbecher M, Raynaud L. 2010. 'Ensemble of data assimilations at ECMWF'. ECMWF Technical Memorandum 636, Reading, UK.
- Jakob C. 2010. Accelerating progress in global atmospheric model development through improved parametrizations. *Bull. Am. Meteorol. Soc.* **91**: 869–875.
- Jung T, Palmer TN, Shutts GJ. 2005. Influence of stochastic parametrization on the frequency of occurrence of North Pacific weather regimes in the ECMWF model. *Geophys. Res. Lett.* **32**: L23811, DOI: 10.1029/2005GL024248.
- Jung T, Miller MJ, Palmer TN, Towers P, Wedi N, Achuthavarier D, Adams JM, Altshuler EL, Cash BA, Kinter JL III, Marx L, Stan C, Hodges KI. 2012. High-resolution global climate simulations with the ECMWF model in Project Athena: experimental design, model climate and seasonal forecast skill. *J. Climate* (in press).
- Khouider B, Majda AJ, Katsoulakis MA. 2003. Coarse-grained stochastic models for tropical convection and climate. *Proc. Natl Acad. Sci. USA* **100**: 11941–11946.
- Kinter JL III, Cash B, Achuthavarier D, Adams J, Altshuler E, Dirmeyer P, Doty B, Huang B, Marx L, Manganello J, Stan C, Wakefield T, Jin E, Palmer T, Hamrud M, Jung T, Miller M, Towers P, Wedi N, Satoh M, Tomita H, Kodama C, Nasuno T, Oouchi K, Taniguchi H, Andrews P, Baer T, Ezell M, Halloy C, John D, Loftis B, Mohr R, Wong K. 2012. Revolutionizing climate modeling – Project Athena: a multi-institutional international collaboration. *Bull. Am. Meteorol. Soc.* (in press).
- Lander J, Hoskins BJ. 1997. Believable scales and parametrizations in a spectral transform model. *Mon. Weather Rev.* **125**: 292–303.
- Lewis JM. 2005. Roots of ensemble forecasting. *Mon. Weather Rev.* **133**: 1865–1885.
- Lilly DK. 1973. Lectures in sub-synoptic scales of motions and two-dimensional turbulence. In *Dynamic Meteorology*, Morel P (ed.). Reidel: Boston, MA; pp. 353–418.
- Lin JW-B, Neelin JD. 2003. Toward stochastic deep convective parameterization in general circulation models. *Geophys. Res. Lett.* **30**: 1162, DOI: 10.1029/2002GL016203.
- Lindborg E. 2007. Horizontal wavenumber spectra of vertical vorticity and horizontal divergence in the upper troposphere and lower stratosphere. *J. Atmos. Sci.* **64**: 1017–1025.
- Lorenz EN. 1969. The predictability of a flow which possesses many scales of motion. *Tellus* **21**: 289–307.
- Lorenz EN. 1975. Climatic predictability. In *The Physical Basis of Climate and Climate Modelling*. WMO GARP Publication Series No 16, World Meteorological Organisation: Geneva.
- Lorenz EN. 1996. Predictability: a problem partly solved. In *Proceedings: Seminar on Predictability*. ECMWF. Reprinted in Palmer TN, Hagedorn R. 2006. *Predictability of Weather and Climate*. Cambridge University Press: Cambridge, UK; 40–48.
- Majda AJ, Bertozzi AL. 2001. *Vorticity and Incompressible Flow*. Cambridge Texts in Applied Mathematics, Cambridge University Press: Cambridge, UK.
- Manabe S, Wetherald RT. 1975. The effects of doubling the CO₂ concentration on the climate of a general circulation model. *J. Atmos. Sci.* **32**: 3–15.
- Masson D, Knutti R. 2011. Climate model genealogy. *Geophys. Res. Lett.* **38**: L08703, DOI: 10.1029/2011GL046864.
- Matsueda M, Palmer TN. 2011. Accuracy of climate change predictions using high resolution simulations as surrogates of truth. *Geophys. Res. Lett.* **38**: L05803, DOI:10.1029/2010GL046618.
- Meehl GA, Covey C, Delworth T, Latif M, McAvaney B, Mitchell JFB, Stouffer RJ, Taylor KE. 2007. The WCRP CMIP3 multimodel dataset. *Bull. Am. Meteorol. Soc.* **42**: 950–960.
- Murphy AH. 1969. Measures of the utility of probabilistic predictions in cost-loss ratio decision situations in which knowledge of the cost-loss ratios is incomplete. *J. Appl. Meteorol.* **8**: 863–873.
- Murphy JM, Palmer TN. 1986. Experimental monthly forecasts for the United Kingdom. Part II: A real-time long-range forecast by an ensemble of numerical integrations. *Meteorol. Mag.* **115**: 337–344.

- Murphy JM, Sexton DMH, Barnett DN, Jones GS, Webb MJ, Collins M, Stainforth DA. 2004. Quantification of modelling uncertainties in a large ensemble of 265 climate change simulations. *Nature* **430**: 768–772.
- Nastrom GD, Gage KS. 1985. A climatology of atmosphere wavenumber spectra of wind and temperature observed by commercial aircraft. *J. Atmos. Sci.* **43**: 857–870.
- Palem KV. 2005. Energy aware computing through probabilistic switching: a study of limits. *IEEE Trans. Comput.* **54**: 1123–1137.
- Palmer TN. 1997. 'On parametrising scales that are only somewhat smaller than the smallest resolved scale, with application to convection and orography'. In *Proceedings of the ECMWF Workshop on New Insights and Approaches to Convective Parametrisation*. ECMWF: Reading, UK; 328–337.
- Palmer TN. 1998. Climate change from a nonlinear dynamical perspective. *J. Climate* **12**: 575–591.
- Palmer TN. 2001. A nonlocal dynamical perspective on model error: a proposal for nonlocal stochastic-dynamic parametrization in weather and climate prediction models. *Q. J. R. Meteorol. Soc.* **127**: 279–304.
- Palmer TN. 2011. A CERN for climate change. *Physics World* **24**: 14–15.
- Palmer TN, Webster PJ. 1993. 'Towards a unified approach to climate and weather prediction'. In *Proceedings of 1st Demetra Conference on Climate Change*. European Community Press: Brussels.
- Palmer TN, Weisheimer A. 2011. Diagnosing the causes of bias in climate models: why is it so hard? *Geophys. Astro. Fluid* **105**: 351–365.
- Palmer TN, Doblas-Reyes FJ, Weisheimer A, Rodwell MJ. 2008. Towards seamless prediction: calibration of climate change projections using seasonal forecasts. *Bull. Am. Meteorol. Soc.* **89**: 459–470.
- Palmer TN, Buizza R, Doblas-Reyes F, Jung T, Leutbecher M, Shutts GJ, Steinheimer M, Weisheimer A. 2009. 'Stochastic parametrization and model uncertainty'. Technical Report 598, European Centre for Medium-Range Weather Forecasts: Reading, UK.
- Pennell C, Reichler T. 2011. On the effective number of climate models. *J. Climate* **24**: 2358–2367.
- Phillips NA. 1956. The general circulation of the atmosphere: a numerical experiment. *Q. J. R. Meteorol. Soc.* **82**: 123–164.
- Plant RS, Craig CG. 2008. A stochastic parameterization for deep convection based on equilibrium statistics. *J. Atmos. Sci.* **65**: 87–105.
- Pollard D. 2002. *A User's Guide to Measure Theoretic Probability*. Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press: Cambridge, UK.
- Rodwell MJ, Palmer TN. 2007. Using numerical weather prediction to assess climate models. *Q. J. R. Meteorol. Soc.* **133**: 129–146.
- Rossow WL, Cairns B. 1995. Monitoring changes of clouds. *Climate Change* **31**: 305–347.
- Rotunno R, Snyder C. 2008. A generalization of Lorenz's model for the predictability of flows with many scales of motion. *J. Atmos. Sci.* **65**: 1063–1076.
- Scaife AA, Spanghel T, Fereday D, Cubasch U, Langematz U, Akiyoshi H, Bekki S, Braesicke P, Butchart N, Chipperfield M, Gattelman A, Hardiman S, Michou M, Rozanov E, Shepherd TG. 2011. Climate change and stratosphere–troposphere interaction. *Clim. Dynam.* DOI 10.1007/s00382-011-1080-7.
- Schertzer D, Lovejoy S. 1993. 'Lecture notes: nonlinear variability in geophysics 3: Scaling and multifractal processes in geophysics. Institut d'Etudes Scientifique de Cargèse, Cargèse, France.
- Schertzer D, Lovejoy S. 2004. Uncertainty and unpredictability in geophysics: chaos and multifractal insights. State of the planet: frontiers and challenges in geophysics. *Geophys. Monogr. Series* **150**: 317–334.
- Shukla J, Palmer TN, Hagedorn R, Hoskins B, Kinter J, Marotzke J, Miller M, Slingo J. 2010. Toward a new generation of world climate research and computing facilities. *Bull. Am. Meteorol. Soc.* **91**: 1407–1412.
- Shutts G. 2005. A kinetic energy backscatter algorithm for use in ensemble prediction systems. *Q. J. R. Meteorol. Soc.* **131**: 3079–3102.
- Shutts GJ, Palmer TN. 2007. Convective forcing fluctuations in a cloud-resolving model: relevance to the stochastic parametrization problem. *J. Climate* **20**: 187–202.
- Simmons AJ, Hollingsworth A. 2002. Some aspects in the improvement in skill in numerical weather prediction. *Q. J. R. Meteorol. Soc.* **128**: 647–677.
- Slingo J, Palmer TN. 2011. Uncertainty in weather and climate prediction. *Phil. Trans. R. Soc. A* **369**: 4751–4767.
- Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds). 2007. *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, 2007*. Cambridge University Press: Cambridge, UK.
- Stainforth D, Aina T, Christensen C, Collins M, Faull N, Frame DJ, Kettleborough JA, Knight S, Martin A, Murphy JM, Piani C, Sexton D, Smith LA, Spicer RA, Thorpe AJ, Allen MR. 2005. Uncertainty in predictions of the climate response to rising levels of greenhouse gases. *Nature* **433**: 403–406.
- Stephens GL. 2005. Cloud feedbacks in the climate system: a critical review. *J. Climate* **18**: 237–273.
- Straus DM, Corti S, Molteni F. 2007. Circulation regimes: chaotic variability versus SST-forced predictability. *J. Climate* **20**: 2251–2272.
- Thuburn J, Kent J, Wood N. 2011. 'Energy and entropy cascades in numerical models'. In *Proceedings of ECMWF Workshop on Representing Model Error in Weather and Climate prediction*.
- Tompkins AM, Berner J. 2008. A stochastic convective approach to account for model uncertainty due to unresolved humidity variability. *J. Geophys. Res.* **113**: D18101.
- WCRP. 2005. 'The World Climate Research Programme Strategic Framework, 2005–2015 Coordinated Observation and Prediction of the Earth System (COPES)'. Technical Document No. 1291 (WCRP-123), World Meteorological Organization: Geneva.
- Webster PJ, Jian J, Hopson TM, Hoyos CD, Agudelo PA, Chang H-R, Curry JA, Grossman RL, Palmer TN, Subbiah AR. 2010. Extended-range probabilistic forecasts of Ganges and Brahmaputra floods. *Bull. Am. Meteorol. Soc.* **91**: 1493–1514.
- Weisheimer A, Doblas-Reyes FJ, Palmer TN, Alessandri A, Arribas A, Deque M, Keenlyside N, MacVean M, Navarra A, Rogel P. 2009. ENSEMBLES: a new multi-model ensemble for seasonal-to-annual predictions: skill and progress beyond DEMETER in forecasting tropical Pacific SSTs. *Geophys. Res. Lett.* **36**: L21711, DOI: 10.1029/2009GL040896.
- Weisheimer A, Palmer TN, Doblas-Reyes FJ. 2011. Assessment of representations of model uncertainty in monthly and seasonal forecast ensembles. *Geophys. Res. Lett.* **38**: L16703, DOI: 10.1029/2011GL048123.
- Williams K, Webb M. 2009. A quantitative performance assessment of cloud regimes in climate models. *Clim. Dynam.* **33**: 141–157.
- Wilks DS. 2005. Effects of stochastic parametrizations in the Lorenz '96 system. *Q. J. R. Meteorol. Soc.* **131**: 389–407.
- Wilks DS. 2006. *Statistical Methods in the Atmospheric Sciences*. Vol. 1 of *International Geophysics Series*, Elsevier: Amsterdam.
- Woollings T, Hoskins BJ, Blackburn M, Berrisford P. 2008. A new Rossby wave-breaking interpretation of the North Atlantic Oscillation. *J. Atmos. Sci.* **65**: 609–626.